

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau(43) International Publication Date  
22 July 2004 (22.07.2004)

PCT

(10) International Publication Number  
WO 2004/060834 A1(51) International Patent Classification<sup>7</sup>: C07C 1/00,  
B01D 17/06(21) International Application Number:  
PCT/IL2003/001112(22) International Filing Date:  
29 December 2003 (29.12.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
153792 2 January 2003 (02.01.2003) IL(71) Applicants (for all designated States except US): YIS-  
SUM RESEARCH DEVELOPMENT COMPANY  
OF THE HEBREW UNIVERSITY OF JERUSALEM  
[IL/IL]; Hi Tech Park, Edmond J Safra Campus, Givat  
Ram, P.O. Box 39135, Jerusalem 91390 (IL). ARKAL  
FILTRATION SYSTEMS C.S. LTD. [IL/IL]; Bet Zera,  
15135 D.N. Emek Hayarden (IL).


(72) Inventors; and

(75) Inventors/Applicants (for US only): JOSHI, Ashutosh

[IN/IN]; B103-4 Rushikes Apts., Kacheri Road, Kartaj,  
410201 Raigad M.S. (IN). KOLODNY, Yuri [IL/IL]; 3/6  
Raphael Elias Street, 67531 Tel Aviv (IL). KREISEL, Itay  
[IL/IL]; 10 Simtat Aharon Street, 34363 Haifa (IL). SAS-  
SON, Yoel [IL/IL]; 30 HaPalmach Street, 92542 Jerusalem  
(IL).(74) Agents: LUZZATTO, Kfir et al.; Luzzatto & Luzzatto,  
P.O. Box 5352, 84152 Beersheva (IL).(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR,  
CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,  
GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,  
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,  
MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU,  
SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,  
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.(84) Designated States (regional): ARIPO patent (BW, GH,  
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,

[Continued on next page]

(54) Title: ENHANCED GENERATION OF HYDROXYL RADICALS



(57) Abstract: This invention provides a process for enhancing the generation of hydroxyl radicals in aqueous mixtures containing hydrogen peroxide, which process comprises supplying oxygen and magnesium oxide to the mixture and irradiating it with UV light. The process can be used, for example, for processing ballast water, industrial waste waters, and municipal waste waters.

WO 2004/060834 A1



SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**Published:**

— *with international search report*

## ENHANCED GENERATION OF HYDROXYL RADICALS

### Field of the Invention


The present invention relates to a method for enhancing the formation of hydroxyl radicals in water environment in the presence of hydrogen peroxide, by the combined effect of UV irradiation, dissolved oxygen, and magnesium oxide as a catalyst.

### Background of the Invention


Hydroxyl radical (HO\*) is one of the strongest oxidants that is available in the natural world and in the chemist's tool box. This species is extremely reactive and degrades promptly any available organic molecule. Hydroxyl radicals, likewise other reactive oxygen species, are also a very potent biocide which vigorously attack microorganisms and tissues, damaging cell membrane lipids, proteins, carbohydrates and DNA. The hydroxyl radical reactions are extremely fast, having rate constants of from  $10^7$  to  $10^9$   $M^{-1} \text{ sec}^{-1}$ . Consequently the hydroxyl radical only diffuses the equivalent of 5-10 of its molecular diameter before it reacts [E. Cadenas: Ann. Rev. Biochem. 58 (1989) 79]. Interestingly, hydroxyl radicals and other reactive oxygen species were recently recognized as a cellular signaling device [E.g., Van Breusegem et al.: Plant Sci. 161 (2001) 405-414], as well as a biological weapon used by the immune system for killing bacteria [Wentworth P. et al.: Science 298 (2002) 2195-9].

Hydroxyl radicals injure cell membrane lipids via the following chain mechanism:


a) Detachment of a hydrogen radical randomly from a fatty acid, leading to the formation of an alkyl radical:



b) The alkyl radical reacts instantly with oxygen (which is available in aerobic environment) to yield an alkylhydroperoxide derivative:




c) The alkyl hydroperoxide reacts with water to generate an alcohol and a hydroperoxyl radical:




5 The transformation of a fatty acid skeleton into an alcohol renders the destruction of the membrane structure and consequently the collapse of the cell wall and the death of the organism.

d) The hydroperoxyl radical absorbs a hydrogen radical from another fatty acid molecule to yield an alkyl radical that subsequently reacts with oxygen,  
10 as shown in step b):




The hydrogen peroxide formed becomes the source of additional hydroxyl radicals which renew the cycle. This chain process propagates until a termination step such as the following one takes place:



A fundamental feature of hydroxyl radicals (and other reactive oxygen species) is the benign nature of their decomposition products. No chemicals are formed as a result of their application in various water treatments but  
20 water.


Several methods for the generation of hydroxyl radicals are known, based mainly on hydrogen peroxide as the source. The simplest and the most accepted methodology is based on the Fenton reaction where ferrous ion  
25 reacts with hydrogen peroxide in an acidic media as follows [Legrini O.: Chem. Rev. 93 (1993) 671-98]:




This reaction is made catalytic predominantly under the effect of UV irradiation of the wavelength greater than 300 nm, which allows the  
30 recycling of  $Fe^{+3}$  to the original  $Fe^{+2}$ , which process is named "Photo-Fenton" reaction. The main drawback of the Fenton type procedures is their strong

dependence on pH. Thus the above reaction occurs only under acidic conditions, preferably at pH= 2.5-3.0.[Andreozzi R. et al.: Catalysis Today 53 (1999) 51-9]. Consequently the Fenton scheme is not practicable under the numerous situations where acidic media cannot be used, such as in ballast  
5 water treatment, for example.



Another procedure for the formation of hydroxyl, and hydroperoxyl, radicals uses photocatalysis together with semi-conductors such as TiO<sub>2</sub>. Upon irradiation of TiO<sub>2</sub> particle by the light having wavelength about 315-395  
10 nm, an electron is ejected, generating a "hole" on the particle surface:




The "hole" interacts with water to produce a hydroxyl radical:



The free electron reacts with oxygen molecule to produce an anion of  
15 superoxide radical which is protonated to yield a hydroperoxyl radical:



Another widespread technique for the making of hydroxyl radical is direct  
20 photolysis of hydrogen peroxide under UV irradiation:



This procedure suffers from low efficacy due to the very low molar extinction coefficient of the hydrogen peroxide molecule (about 19 dm<sup>3</sup>.mol<sup>-1</sup> .cm<sup>-1</sup> at 254 nm at neutral pH [Baxendale J.H.: Trans. Faraday Soc. 53 (1957) 344].

25

Using EPR techniques, Giamello E. et al. [J. Phys. Chem. 97(1993) 5735-40] showed that reactive oxygen species were formed upon contact of hydrogen peroxide with the surface of magnesium oxide in an aqueous environment. These authors also observed a highly unexpected stability of hydroxyl  
30 radicals adsorbed on the MgO surface, wherein the radicals were stable at temperatures as high as 200° C.

It is an object of this invention to provide a high efficacy method for producing hydroxyl radicals in aqueous environment using UV irradiation of hydrogen peroxide solutions.

- 5 It is another object of this invention to provide a method for producing hydroxyl radicals in water solutions of hydrogen peroxide and oxygen, combining UV irradiation with a catalytic effect of magnesium oxide.

10 Other objects and advantages of present invention will appear as description proceeds.

### **Summary of the Invention**

This invention provides a method for enhancing the generation of hydroxyl radicals in aqueous mixtures containing hydrogen peroxide comprising: i) supplying oxygen or air to said mixture; ii) supplying magnesium oxide to said mixture; iii) irradiating said mixture with UV light, preferably having a wavelength of from 190 to 390 nm; and iv) mixing said mixture. Said aqueous mixture is an aqueous solution or suspension. The initial concentration of hydrogen peroxide is from 2 ppm to 250 ppm, preferably 10 ppm to 50 ppm. Magnesium oxide is preferably added to the mixture in one portion of from 2 ppm to 250 ppm, and more preferably from 10 ppm to 50 ppm. Oxygen is supplied as air or pure oxygen, preferably to saturation.

### **Brief Description of the Drawings**

25 The above and other characteristics and advantages of the invention will be more readily apparent through the following examples, and with reference to the appended drawings, wherein:

Fig. 1. Shows the effect of individual factors on the formation of hydroxyl radicals;

30 Fig. 2. is a HPLC chromatogram of salicylic and hydroxysalicylic acid; and

Fig. 3. is a table summarizing conversion rates of salicylic acid under various conditions.

### Detailed Description of the Invention

5 It has now been surprisingly found that the presence of magnesium oxide in an aqueous mixture containing hydrogen peroxide and oxygen leads to enhanced formation of hydroxyl radicals under UV irradiation. Without wishing to be limited by any particular theory, the inventors believe that magnesium oxide is a strong catalyst during the creation of hydroxyl radicals  
10 in water environment provided the system is irradiated by a UV light. This phenomenon provides particularly good results in the presence of oxygen. The synergic combination of hydrogen peroxide, oxygen, magnesium oxide and UV irradiation provides an effective new means for the generation of hydroxyl radicals in water. Surprisingly, even in the absence of hydrogen  
15 peroxide, a significant amount of hydroxyl radicals are formed in water in the presence of oxygen and magnesium oxide. In a preferred embodiment of the present invention, hydrogen peroxide is added to an aqueous solution in which the formation of hydroxyl radicals is desired, and magnesium oxide is injected as a suspension while mixing the aqueous solution, introducing into  
20 it oxygen, and irradiating it by UV. Any commercially available UV irradiation source can be used.

In a preferred embodiment of the method according to this invention, the initial concentration of hydrogen peroxide is from 2 ppm to 250 ppm, the  
25 solution is saturated with oxygen, and magnesium oxide in suspension is injected as a catalyst to a concentration from 2 ppm to 250 ppm. Oxygen is preferably introduced into the mixture by injection of oxygen or air.

In one of the preferred embodiments of the invention, the peroxide radicals  
30 serve for the treatment of ballast water. In a preferred embodiment of this invention, hydrogen peroxide is added to sea water, intended to serve as

ballast water, to a concentration of from 10 to 50 ppm, air or oxygen is finely dispersed to saturate the mixture, and magnesium oxide is added to a concentration of from 10 ppm to 50 ppm.

5 This invention thus provides a method for enhancing the generation of hydroxyl radicals in aqueous mixtures containing hydrogen peroxide comprising i) supplying oxygen to said mixture; ii) supplying magnesium oxide to said mixture; iii) irradiating said mixture with UV light; and iv) mixing said mixture for a period sufficient to generate the desired amount of radicals, wherein said desired amount of radicals may be an amount  
10 sufficient to reach a required effect in the mixture, for example a biocidal effect. Said desired amount of radicals may be a required predetermined quantity. Hydroxyl radicals may be quantified by physical or chemical methods known in the art, such as measuring redox potentials, or reacting hydroxyl radicals with a trapping agent such as salicylic acid, etc. Said  
15 aqueous mixture can be a solution or suspension. Air or oxygen is supplied in any suitable manner that can inject and homogenize it into the aqueous solution, preferably providing saturation. According to a preferred embodiment of the invention the UV radiation has a wavelength of from 190  
20 to 390 nm. Magnesium oxide is preferably added to the mixture in one portion. The period that enables the generation of required amount of radicals depends on the type of application of said mixture, and may be typically from few seconds to several hours, for example from 3 seconds to 5 hours, and preferably it is from 30 seconds to 100 minutes. The mixture may  
25 be irradiated essentially during the whole period that is set for the generation of hydroxyl radicals, and also oxygen may be supplied to the mixture during the whole of said period, or, alternatively, the application of UV and supplying oxygen may last only a few seconds, if the required effect is attained that quickly in said type of application.



In a preferred embodiment of this invention, the pH of the aqueous solution, in which the formation of hydroxyl radicals is desired, has a value of from 5 to 10, and preferably from 7.2 to 9.7.

5 The described process can be used whenever the increased concentration of hydroxyl radicals is desirable, an example being processing ballast water, industrial waste waters, municipal waste waters, etc. The method of this invention may be thus used, for example, for the purification of sea water or municipal effluent water, however without being limited to these examples.

10


The invention will be further described and illustrated by the following examples.

## Examples

15

### Quantifying hydroxyl radicals

The formation of hydroxyl radicals was quantified by reacting the radicals with salicylic acid [H. Kaur et al.: Methods in Enzymology 233 (1994) 67-82] to form 2,3-dihydroxybenzoic acid and 2,5-dihydroxybenzoic acid, and by detecting these products by HPLC with a UV detector (See Fig. 2), using  
20 column RP-18, 80% phosphate buffer pH 6.5 + 20% methanol, flow rate 1 ml/min, and  $\lambda_{\max}$  300 nm.



25

The degree and rate of the conversion to dihydroxybenzoic acid is proportional to the amount of hydroxyl radicals generated in a given system.

### Example 1

Reaction was carried out in a one liter beaker equipped with a stirrer, into which a medium pressure mercury UV lamp in a quartz sleeve, Immersion Lamp TQ 150W, was immersed. Approximately half the lamp was immersed.

5 Tap water passed continuously through quartz jacket to cool it. The total volume of reaction mixture was 800 ml. Illuminated length was 41 mm. Salicylic acid (Fluka, purity > 99.5 %), 500 ppm in 400 ml seawater, and 400 ml seawater were stirred with a magnetic stirrer, and the pH of the solution was adjusted to 7.85 with 0.5 % NaOH. The irradiation continued for 90

10 min. Samples were withdrawn at 10, 20, 40, 60 and 90 min, and analyzed by HPLC. The conversion rates were calculated as the ratio of (OH-salicyl.)/(salicyl. + OH-salicyl.) (see Figs. 1, 2, and 3).

### Example 2

15 The reaction was started as in Example 1, but 20 ppm (16 mg) of magnesium oxide (obtained from Aldrich, purity 98%) was added to 800 ml of the reaction mixture after the pH adjustment, and after 5 min the irradiation started, and continued for 90 min.

### Example 3

20 The reaction was started as in Example 1, but hydrogen peroxide, 20 ppm in 400 ml seawater, was added to salicylic acid, 500 ppm in 400 ml sea water, pH was adjusted to 7.85 with 0.5 % NaOH, and the mixture was then irradiated during 90 min.

### Example 4

25 The reaction was started as in Example 3, but 20 ppm (16 mg) of magnesium oxide was added to 800 ml of the reaction mixture after the pH adjustment, and only then the irradiation started.

### Example 5

30 The reaction was started as in Example 1, but after the pH adjustment, purging of the mixture started, oxygen being fed at a rate of 50 ml/min

through a glass tube of diameter 0.5 cm. After 10 min, the irradiation started, and continued for 90 min.

#### Example 6

5 The reaction was started as in Example 2, but after the addition of magnesium oxide, and 5 min stirring, purging of the mixture with oxygen started, oxygen being introduced through a glass tube of diameter 0.5 cm. After 10 min, the irradiation started, and continued for 90 min.

#### Example 7

10 The reaction was started as in Example 3, but after the pH adjustment, the mixture was purged with oxygen which was introduced through a glass tube of diameter 0.5 cm. After 10 min, the irradiation started, and continued for 90 min.

#### Example 8

15 The reaction was started as in Example 4, but after adding magnesium oxide and 5 min stirring, purging of the mixture with oxygen started, oxygen being introduced through a glass tube of diameter 0.5 cm. After 10 min, the irradiation started, and continued for 90 min. The pH at 90 min was 7.91.

20 The initial and final temperature of the reaction mixture were 23°C and 25.5°C, respectively, and the initial and final redox potential of the reaction mixture were 180 mV and 110 mV, respectively. The superior generation of hydroxyl radicals in the system  $\text{MgO}/\text{H}_2\text{O}_2/\text{O}_2$  can be seen in Figs. 1 and 3.

#### Example 9

25 Some of the enhanced activity, seen in the mixtures containing MgO, might be attributed to the increased basicity in the presence of solid MgO. Therefore, two reactions were performed comparing the effect of MgO with the effect of sodium hydroxide. Two mixtures contained 250 ppm salicylic acid, 30 ppm hydrogen peroxide, and they were irradiated by UV. The first

30 mixture further contained 250 ppm MgO, its pH was adjusted to 7.8 at the

beginning of the reaction, and rose to 9.38 after 90 minutes. The second mixture did not contain MgO and its initial pH was adjusted to 9.55 by NaOH. Magnesium oxide had stronger effect even when compared to the effect of sodium hydroxide, pH 9.55, which further demonstrates the phenomenon of catalytic properties of MgO.

While the invention has been described using some specific examples, many modifications and variations are possible. It is therefore understood that the invention is not intended to be limited in any way, other than by the scope of the appended claims.

## CLAIMS

1. A method for enhancing the generation of hydroxyl radicals ( $\text{OH}^*$ ) in aqueous mixtures containing hydrogen peroxide, comprising
  - i) supplying oxygen to said mixture;
  - ii) supplying magnesium oxide to said mixture as a catalyst;
  - iii) irradiating said mixture with UV light; and
  - iv) mixing said mixture.
2. The method of claim 1, wherein the aqueous mixture is an aqueous solution or suspension.
3. The method of claim 1, wherein hydrogen peroxide has an initial concentration of from 2 to 250 ppm.
4. The method of claim 1, wherein oxygen is supplied by injecting of air or oxygen into the mixture.
5. The method of claim 1, wherein oxygen is supplied to saturation.
6. The method of claim 1, wherein said UV radiation has wavelength of from 190 to 390 nm.
7. The method of claim 1, wherein magnesium oxide is added to the mixture to a concentration of from 2 ppm to 250 ppm.
8. The method of any one of claims 1 to 7, wherein the initial concentration of hydrogen peroxide is from 10 to 50 ppm, and the initial concentration of magnesium oxide is from 10 to 50 ppm.
9. The method of any one of claims 1 to 8, wherein the pH of said mixture has a value of from 5 to 10.

10. The method of claim 9, wherein said pH has a value of 7.2 to 9.7.
11. The method of claim 1, wherein said mixing is carried out for a period of time sufficient to generate the desired amount of radicals.
12. The method of claim 11, wherein said desired amount of radicals is an amount sufficient to reach a required biocidal effect in the mixture.
13. The method of claim 11, wherein said period lasts from 3 seconds to 5 hours.
14. The method of claim 13, wherein said period lasts from 30 second to 100 minutes.
15. The method of claim 11, wherein said period lasts more than 5 hours.
16. The method of claim 11, wherein said desired amount of radicals is a predetermined quantity.
17. The method of claim 11, wherein generated radicals are quantified by a physical or chemical method.
18. The method of claim 17, wherein said chemical method comprises reacting hydroxyl radicals with salicylic acid.

1/3




Fig. 1

2/3




Fig. 2



3/3

Run No	React. cond.	H <sub>2</sub> O <sub>2</sub> (ppm)	MgO (ppm)	Conv. of SA in 90 min (%)	Conv. of SA in 20 min (%)
1	SA/UV	0	0	8.42	5.8
2	SA/UV/MgO	0	20	10.51	6.12
3	SA/UV/H <sub>2</sub> O <sub>2</sub>	10	0	11.56	6.64
4	SA/UV/H <sub>2</sub> O <sub>2</sub> /MgO	10	20	15.42	8.2
5	SA/UV/ O <sub>2</sub>	0	0	15.08	5.9
6	SA/UV/MgO/ O <sub>2</sub>	0	20	20.56	6.89
7	SA/UV/H <sub>2</sub> O <sub>2</sub> /O <sub>2</sub>	10	0	20.31	8.26
8	SA/UV/H <sub>2</sub> O <sub>2</sub> /MgO /O <sub>2</sub>	10	20	27.37	11.76

Fig. 3

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/IL03/01112

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07C 1/00; B01D 17/06

US CL : 204/157.15; 210/748

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 204/157.15; 210/748

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
Please See Continuation Sheet

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	PANKRATOV, A., Photodissociation of Water, Heavy Water and Hydrogen Peroxide on a Magnesium Oxide Surface, Khimiya Vysokikh Energii, (no month) 1970, Vol. 4, No. 2, pp. 126-130, abstract.	1-18



Further documents are listed in the continuation of Box C.



See patent family annex.

\* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "B" earlier application or patent published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

20 April 2004 (20.04.2004)

Date of mailing of the international search report

06 MAY 2004

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US  
Commissioner for Patents  
P.O. Box 1450  
Alexandria, Virginia 22313-1450

Facsimile No. (703) 305-3230

Authorized officer

Edna Wong

Telephone No. (572) 272-1700

*Edna Wong*

**INTERNATIONAL SEARCH REPORT**

PCT/IL03/01112

**Continuation of B. FIELDS SEARCHED Item 3:**

**EAST, STN**

search terms: hydroxyl radicals, hydrogen peroxide, magnesium oxide, UV